

Table 36. Deaths due to coronary heart disease among San Diego County Residents, 1995-1998, 1999*

By Region, Gender, Race/Ethnicity, and Age Group

Variable	1995		1996		1997		1998		1999	
	Deaths	Adjusted Rate	Deaths	Adjusted Rate	Deaths	Adjusted Rate	Deaths	Adjusted Rate	Deaths	Adjusted Rate
San Diego County	4,636	203.3	4,562	194.2	4,608	189.9	4,691	186.7	4,676	179.2
Region										
N. Coastal	712	182.7	703	174.6	698	166.4	741	170.2	707	155.7
N. Central	815	190.7	810	183.3	871	188.3	864	176.7	862	169.3
Central	798	236.7	739	218.0	748	215.9	783	224.1	816	230.7
South	632	234.6	603	219.1	597	211.5	639	215.1	627	203.8
East	901	235.1	868	220.4	897	221.3	870	209.0	869	202.8
N. Inland	752	161.6	810	167.1	770	154.5	773	149.2	767	141.5
Gender										
Male	2,345	253.9	2,265	236.8	2,266	230.1	2,405	233.5	2,409	226.1
Female	2,291	163.9	2,297	159.1	2,342	157.1	2,286	149.2	2,267	142.1
Race/Ethnicity										
White	3,953	209.8	3,864	200.5	3,906	197.6	3,922	193.5	3,880	185.7
Black	171	307.3	185	301.0	181	294.5	187	292.2	192	274.6
Hispanic	351	151.2	342	141.3	331	130.6	373	133.1	386	129.5
Asian/Other	161	136.7	171	135.5	190	141.0	207	136.6	217	136.0
Age Groups										
Under 1	0	0.0	0	0.0	0	0.0	0	0.0	1	*
1-4	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
5-14	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
15-24	1	*	4	*	0	0.0	1	*	2	*
25-34	11	2.4	10	2.2	10	2.3	6	1.4	11	2.5
35-44	50	11.4	43	9.7	37	8.2	58	12.6	61	13.1
45-54	160	56.2	180	60.6	170	54.6	172	52.4	175	50.9
55-64	361	195.7	341	180.5	358	183.0	405	197.7	361	168.3
65-74	955	574.9	869	525.2	837	503.0	830	491.2	800	465.6
75-84	1,510	1,459.1	1,547	1,461.4	1,519	1,402.4	1,590	1,435.6	1,504	1,331.8
85+	1,588	4,712.2	1,568	4,388.8	1,677	4,437.2	1,629	4,081.6	1,761	4,168.8

All rates are per 100,000 population and are adjusted to 2000 Standard U.S. Population.

* No rates calculated for fewer than 5 events.

Note: Rates based on fewer than 20 events should be interpreted with caution.

*Note: The 1999 mortality data is the first release to use ICD-10 to code underlying cause of death and differs from the ICD-9 classification used for the 1995-1998 data. Due to lack of comparability between the two ICD versions, the differences in deaths and rates between pre-1999 and 1999 may result in major discontinuities in trend data.

Source: State of California, Department of Health Services, Death Statistical Master Files, 1995-1999; SANDAG, January 1, 1995 -1999 population estimates used for calculating rates.