

Table 42. Deaths due to prostate cancer in males among San Diego County Residents, 1995-1998, 1999*

By Region, Race/Ethnicity, and Age Group

| Variable | 1995 | | 1996 | | 1997 | | 1998 | | 1999 | |
|-------------------------|---------------|--------------------------|---------------|--------------------------|---------------|--------------------------|---------------|--------------------------|---------------|--------------------------|
| | Deaths | Adjusted Rate | Deaths | Adjusted Rate | Deaths | Adjusted Rate | Deaths | Adjusted Rate | Deaths | Adjusted Rate |
| San Diego County | 283 | 30.9 | 287 | 30.6 | 294 | 30.7 | 266 | 26.7 | 297 | 28.6 |
| Region | | | | | | | | | | |
| N. Coastal | 68 | 42.2 | 60 | 35.1 | 52 | 30.7 | 49 | 27.4 | 43 | 22.9 |
| N. Central | 48 | 26.9 | 51 | 29.5 | 63 | 36.3 | 58 | 29.7 | 58 | 27.8 |
| Central | 34 | 26.5 | 29 | 23.2 | 36 | 27.7 | 33 | 25.4 | 42 | 32.3 |
| South | 31 | 27.4 | 28 | 26.3 | 30 | 25.9 | 28 | 25.4 | 42 | 34.7 |
| East | 47 | 32.0 | 55 | 36.0 | 51 | 32.5 | 42 | 26.2 | 42 | 26.4 |
| N. Inland | 55 | 29.1 | 62 | 30.9 | 60 | 28.9 | 53 | 24.7 | 68 | 30.3 |
| Race/Ethnicity | | | | | | | | | | |
| White | 256 | 33.8 | 242 | 31.5 | 256 | 32.8 | 228 | 28.2 | 265 | 31.6 |
| Black | 10 | 52.5 | 15 | 59.4 | 13 | 55.2 | 8 | 35.4 | 13 | 49.6 |
| Hispanic | 10 | 11.0 | 22 | 21.6 | 17 | 17.2 | 21 | 19.5 | 11 | 9.7 |
| Asian/Other | 7 | 15.8 | 8 | 16.4 | 8 | 14.2 | 9 | 16.7 | 8 | 13.9 |
| Age Groups | Deaths | Age-Specific Rate | Deaths | Age-Specific Rate | Deaths | Age-Specific Rate | Deaths | Age-Specific Rate | Deaths | Age-Specific Rate |
| Under 1 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 |
| 1-4 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 |
| 5-14 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 |
| 15-24 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 |
| 25-34 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 |
| 35-44 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 |
| 45-54 | 5 | 3.5 | 4 | * | 2 | * | 3 | * | 4 | * |
| 55-64 | 14 | 15.8 | 18 | 19.8 | 18 | 19.0 | 12 | 12.1 | 13 | 12.5 |
| 65-74 | 75 | 100.4 | 76 | 101.9 | 59 | 78.4 | 58 | 75.6 | 60 | 76.6 |
| 75-84 | 126 | 298.5 | 119 | 274.9 | 132 | 296.9 | 114 | 249.5 | 143 | 305.8 |
| 85+ | 63 | 587.8 | 70 | 609.0 | 83 | 671.2 | 79 | 594.9 | 77 | 542.1 |

Deaths are for males only. All rates are per 100,000 male population and are adjusted to 2000 Standard U.S. Population.

* No rates calculated for fewer than 5 events.

Note: Rates based on fewer than 20 events should be interpreted with caution.

*Note: The 1999 mortality data is the first release to use ICD-10 to code underlying cause of death and differs from the ICD-9 classification used for the 1995-1998 data. Due to lack of comparability between the two ICD versions, the differences in deaths and rates between pre-1999 and 1999 may result in major discontinuities in trend data.

Source: State of California, Department of Health Services, Death Statistical Master Files, 1995-1999; SANDAG, January 1, 1995-1999 population estimates used for calculating rates.