

Comparison of National, State and County Data for Tuberculosis, 1998-2003

Annual Case Counts
(Rate per 100,000 in parentheses)

	1998	1999	2000	2001	2002	2003
USA	18,371 (6.8)	17,528 (6.4)	16,377 (5.8)	15,991 (5.6)	15,078 (5.2)	14,871 (5.1) ^a
CA	3,855 (11.5)	3,599 (10.9)	3,297 (9.7)	3,332 (9.7)	3,169 (9.0)	3230 (9.1)
San Diego	342 (12.2)	299 (10.4)	295 (10.5)	332 (11.8)	326 (11.2)*	316 (10.7)

* Case rate revised from 10.7 to 11.2 based on revised population estimates for San Diego County for 2002.

^aData Source: Provisional data; National Tuberculosis Surveillance System, DTBE, CDC, March 3, 2004

SAN DIEGO TB STATISTICS, 1997-2003

Table 1. Tuberculosis Incidence by Major Site of Disease, 1997 - 2003

DISEASE SITE	1998	1999	2000	2001	2002	2003
Pulmonary	270 (79%)	219 (73%)	238 (81%)	269 (81%)	246 (75%)	240 (76%)
All Lymphatic	29 (9%)	34 (11%)	30 (10%)	36 (11%)	37 (11%)	34 (11%)
Pleural	9 (3%)	13 (4%)	8 (3%)	6 (2%)	14 (4%)	13 (4%)
Bone/Joint	11 (3%)	8 (3%)	3 (1%)	6 (2%)	9 (3%)	8 (3%)
Peritoneal	3 (1%)	3 (1%)	6 (2%)	2 (<1%)	2 (<1%)	6 (2%)
Meningeal	4 (1%)	5 (2%)	3 (1%)	4 (1%)	2 (<1%)	6 (2%)
Genitourinary	3 (1%)	4 (1%)	3 (1%)	3 (1%)	9 (3%)	2 (<1%)
Miliary/Dissem.	3 (1%)	3 (1%)	0	1 (<1%)	2 (<1%)	1 (<1%)
Other	9 (3%)	8 (3%)	4 (1%)	5 (2%)	5 (2%)	6 (2%)
TOTAL	342	299	295	332	326	316

Table 2. Annual Tuberculosis Cases by Sex, Age and Race/Ethnicity, 1997 - 2003

(Rate per 100,000 in parentheses)

	1998	1999	2000	2001	2002	2003
SEX						
Male	220 (15.4)	173 (11.8)	182 (12.9)	205 (14.4)	195 (12.8)	187 (12.5.)
Female	122 (8.8)	126 (8.9)	113 (8.1)	127 (9.1)	131 (8.8)	129 (8.8)
AGE						
0 - 4	21 (8.5)	18 (7.7)	14 (7.0)	18 (9.1)	10 (4.3)	15 (7.4)
5 - 14	21 (4.9)	23 (5.5)	14 (3.4)	16 (3.9)	4 (0.9)	11 (2.6)
15 - 24	35 (8.8)	24 (5.9)	36 (8.4)	45 (10.4)	39 (8.9)	43 (9.3)
25 - 44	116 (12.8)	99 (10.8)	106 (11.8)	102 (11.3)	130 (13.8)	104 (11.2)
45 - 64	81 (15.2)	67 (11.6)	69 (12.4)	90 (16.1)	81 (12.6)	100 (16)
65 +	68 (21.2)	68 (20.8)	56 (17.8)	61 (19.4)	62 (18.2)	43 (13.2)
RACE/ETHNICITY						
Hispanic	171 (25.3)	138 (19.7)	141 (18.8)	164 (21.8)	154 (19.7)	159 (18.9)
Black	16 (9.5)	22 (12.9)	23 (14.9)	20 (12.9)	27 (15.1)	24 (14.7)
Asian	109 (42.4)	102 (36.8)	103 (40.0)	98 (38.1)	99 (32.9)	89 (32.7)
White	43 (2.9)	34 (1.9)	28 (1.8)	49 (3.2)	46 (2.6)	41 (2.6)
Native Am./Alaskan	3 (--)	3 (--)	0 (--)	1 (--)	0 (--)	0 (--)
Native Hawaiian/ Pacific Islander	--	--	--	--	--	2 (14.4)
Unknown						1 (--)
TOTAL	342 (12.2)	299 (10.4)	295 (10.5)	332 (11.8)	326 (11.2)	316 (10.7)

-- = Data not available

Table 3. Annual Tuberculosis Cases by Country of Origin, 1997 - 2003

	1998	1999	2000	2001	2002	2003
COUNTRY OF ORIGIN						
THE AMERICAS	226	191	179	222	203	219
United States	113	97	72	104	102	107
Mexico	108	88	103	115	96	104
Other	5	7	4	3	5	8
ASIA	105	95	96	96	105	87
Vietnam	14	18	21	18	16	16
Philippines	67	62	62	55	62	45
Other	24	15	13	23	27	26
EUROPE	4	4	2	6	3	1
AFRICA	6	8	14	7	15	9
OTHER/UNKNOWN	1	0	4	1	0	0
TOTAL	342	299	295	332	326	316
% Foreign-born	67	67	76	69	69	66