

COUNTY OF SAN DIEGO

NEWS RELEASE

FOR IMMEDIATE RELEASE Feb. 20, 2009

CONTACT: Leslie Ridgeway (619) 685-2239 School Contact: Anthony Millican (619) 425-9600 ext. 1328

CHICKEN POX CASES REPORTED AT SCHOOL IN CHULA VISTA Five Children Sick; HHSA Urges Parents to Get Their Children Vaccinated

The San Diego County Health and Human Services Agency (HHSA) reports that five students at Kellogg Elementary School in Chula Vista have been diagnosed with chicken pox.

Three of the students had one dose of vaccine; two had no vaccine but had chicken pox before. Two doses of vaccine are recommended to guard against contracting chicken pox, or lessen its effects.

"We recommend that parents talk to their children's healthcare provider about age-appropriate vaccinations," said Wilma Wooten, M.D., M.P.H., County Public Health Officer. "Diseases like chicken pox, whooping cough and diphtheria can cause complications, especially in younger children."

None of the students required hospitalization. School officials have informed staff and parents of students who may have been exposed, and recommended that children have the appropriate vaccinations against the disease.

Chicken pox is a highly contagious disease caused by the varicella virus. The disease is easily spread by coughing, sneezing or contact with chicken pox blisters. The Centers for Disease Control and Prevention recommends two doses of chicken pox vaccine for those who do not have immunity -- the first at 12-15 months of age and the second at 4-6 years of age. A second dose is recommended for individuals who are older and have only had one vaccination,

Symptoms of chicken pox include a skin rash of blister-like lesions covering the body, but usually more concentrated on the face, scalp, and trunk. The risk of complications increases after puberty and includes bacterial infection of skin lesions, dehydration and pneumonia.

Most, but not all, infected individuals have fever, which develops just before or when the rash appears. If exposed, persons who have been vaccinated against the disease may get a milder illness, with less severe rash (sometimes involving only a few red bumps that look similar to insect bites) and mild or no fever. The incubation period is 10-21 days. The illness lasts about 5-10 days. For more information on chicken pox and immunizations in general, please call the HHSA Immunization Branch at 619-692-8661 or visit the website at www.sdiz.org.

###