



COUNTY OF SAN DIEGO

NEWS RELEASE

FOR IMMEDIATE RELEASE

Dec. 4, 2007

CONTACT: Leslie Ridgeway (619) 685-2239

CHICKEN POX OUTBREAK REPORTED AT FUERTE ELEMENTARY SCHOOL *Five Students Ill; Parents Urged to Ensure Their Children Are Immunized*

The San Diego County HHS (Health and Human Services Agency) reports that five students at Fuerte Elementary School in the Cajon Valley Union School District have been diagnosed with chicken pox (varicella). All five students had only one dose of chicken pox vaccine, a reminder to parents that two doses of vaccine is recommended.

"We have worked closely with school officials to identify children who may have been exposed," said Wilma Wooten, M.D., M.P.H., County Public Health Officer. "We urge parents to check with their physicians to make sure their children get two doses of chicken pox vaccine, and are inoculated against other preventable diseases."

Chicken pox is a highly contagious disease caused by the varicella virus. The disease is easily spread by coughing, sneezing or contact with chicken pox blisters. The Centers for Disease Control and Prevention recommends two doses of chicken pox vaccine for all people one year of age and older who do not have evidence of immunity for the disease. Ninety-nine percent of people develop immunity after two doses.

None of the students, all third graders, required hospitalization. School officials have informed parents about the outbreak and asked them to make sure their children have had the appropriate vaccinations against the disease.

Symptoms of chicken pox include a skin rash of blister-like lesions, covering the body but usually more concentrated on the face, scalp, and trunk. The risk of complications increases after puberty and includes bacterial infection of skin lesions, dehydration and pneumonia.

Most, but not all, infected individuals have fever, which develops just before or when the rash appears. If exposed, persons who have been vaccinated against the disease may get a milder illness, with less severe rash (sometimes involving only a few red bumps that look similar to insect bites) and mild or no fever. The incubation period is from 14-16 days from exposure, with a range of 10-21 days. The illness lasts about 5-10 days. For more information on chicken pox and immunizations in general, please call the HHS Immunization Branch at 619-692-8661 or visit the website at www.sdiz.org.

###