

STD Fact Sheet
Sexually Transmitted Diseases by Year of Report
San Diego County, 1993-2004

Year	Syphilis, Infectious (Primary and Secondary)		Congenital Syphilis	
	Cases	Rate*	Cases	Rate†
1993	106	4.0	36	74.0
1994	98	3.6	22	47.0
1995	52	1.9	12	26.4
1996	35	1.3	8	17.6
1997	23	0.8	13	30.1
1998	24	0.9	11	25.4
1999	25	0.9	14	32.4
2000	27	0.9	3	6.8
2001	26	0.9	4	9.2
2002	38	1.3	2	4.6
2003	109	3.7	8	17.8
2004	136	4.5	6	13.1

Year	Gonorrhea		Chlamydia	
	Cases	Rate*	Cases	Rate*
1993	3579	135.2	7720	291.7
1994	2656	98.8	6413	238.6
1995	2176	80.0	5250	193.0
1996	1815	66.7	5642	207.4
1997	1509	55.4	6398	234.8
1998	1595	57.1	7044	252.0
1999	1561	54.7	7591	266.0
2000	1797	61.7	8637	296.6
2001	1875	64.4	9168	314.9
2002	2128	73.1	10225	351.2
2003	1972	66.6	10249	346.1
2004	2376	78.7	10822	358.7

* Rate per 100,000 population

† Rate per 100,000 live births

Sexually Transmitted Diseases
By Age, Gender, Race/Ethnicity and HHS Region
San Diego County, 2004

Characteristic	Syphilis (Primary & Secondary)		Gonorrhea		Chlamydia	
	Cases	Rate*	Cases†	Rate*†	Cases†	Rate*†
<i>AGE GROUP</i>						
< 10	0	0.0	4	1.0	9	2.2
10-14	0	0.0	29	13.1	116	52.3
15-19	2	0.9	402	178.9	2853	1269.3
20-24	10	4.2	638	265.9	4154	1731.6
25-29	21	9.0	419	179.5	1879	804.9
30-34	26	11.0	284	119.9	839	354.2
35-44	48	10.3	424	91.1	725	155.8
≥ 45	29	3.0	176	17.9	247	25.1
<i>GENDER</i>						
Male	126	8.4	1438	95.6	2881	191.5
Female	10	0.7	938	62.0	7941	525.0
<i>RACE/ETHNICITY</i>						
White	82	5.2	855	54.4	3048	193.9
Hispanic	36	4.2	666	77.6	4742	552.6
African Am.	13	8.1	722	449.9	1957	1219.6
Asian/PI	4	1.3	121	39.0	976	314.3
Am. Indian/Alaskan Native	1	6.4	12	76.6	99	631.6
<i>REGION</i>						
North Coastal	7	1.4	141	27.4	1160	225.2
North Inland	2	0.4	131	25.1	1202	230.6
North Central	12	2.1	354	60.9	1815	312.3
Central	98	20.0	1064	217.2	3031	618.8
East	10	2.2	391	84.7	1704	368.9
South	7	1.6	295	67.2	1910	435.0
<i>TOTAL</i>	136	4.5	2376	78.7	10,822	358.7

* Rate per 100,000 population

† Annual estimated cases and rates by age, gender, race/ethnicity and region are obtained by applying the proportion in the known cases to the total case count.