

STD Fact Sheet
Sexually Transmitted Diseases by Year of Report
San Diego County, 1993-2005

Year	Syphilis, Infectious (Primary and Secondary)		Congenital Syphilis	
	Cases	Rate*	Cases	Rate†
1993	106	4.0	36	74.0
1994	98	3.6	22	47.0
1995	52	1.9	12	26.4
1996	35	1.3	8	17.6
1997	23	0.8	13	30.1
1998	23	0.9	11	25.4
1999	25	0.9	14	32.4
2000	27	0.9	3	6.8
2001	27	0.9	4	9.2
2002	38	1.3	2	4.6
2003	109	3.7	8	17.8
2004	136	4.5	6	13.1
2005	193	6.4	4	8.7

Year	Gonorrhea		Chlamydia	
	Cases	Rate*	Cases	Rate*
1993	3579	135.2	7720	291.7
1994	2656	98.8	6413	238.6
1995	2176	80.0	5250	193.0
1996	1815	66.7	5642	207.4
1997	1509	55.4	6398	234.8
1998	1595	57.1	7044	252.0
1999	1561	54.7	7591	266.0
2000	1797	61.7	8637	296.6
2001	1875	64.4	9168	314.9
2002	2128	73.1	10225	351.2
2003	1972	66.6	10249	346.1
2004	2376	78.7	10822	358.7
2005	2606	86.4	11001	364.9

* Rate per 100,000 population

† Rate per 100,000 live births

Sexually Transmitted Diseases
By Age, Gender, Race/Ethnicity and HHS Region
San Diego County, 2005

Characteristic	Syphilis (Primary & Secondary)		Gonorrhea		Chlamydia	
	Cases	Rate*	Cases†	Rate*†	Cases†	Rate*†
AGE GROUP						
< 10	0	0.0	0	0.0	4	1.0
10-14	0	0.0	21	9.7	82	37.7
15-19	8	3.5	513	227.4	2951	1308.3
20-24	18	7.6	699	295.5	4101	1733.8
25-29	28	12.3	485	213.4	2030	893.3
30-34	35	15.1	274	118.5	851	368.1
35-44	68	14.8	439	95.5	728	158.3
≥ 45	36	3.6	175	17.5	254	25.4
GENDER						
Male	169	11.2	1510	100.5	3008	200.1
Female	24	1.6	1096	72.5	7993	528.7
RACE/ETHNICITY						
White	111	7.2	838	54.0	3090	199.2
Hispanic	56	6.4	756	86.5	4932	564.6
African Am.	18	11.3	814	509.7	1961	1227.9
Asian/PI	8	2.6	169	54.2	930	298.4
Am. Indian/Alaskan Native	0	0.0	29	186.3	88	565.2
REGION						
North Coastal	4	0.8	177	36.4	1368	281.5
North Inland	4	0.8	200	39.0	1270	247.6
North Central	21	3.4	406	66.7	1954	320.9
Central	118	24.0	1057	215.1	2961	602.7
East	20	4.4	409	89.3	1535	335.0
South	19	4.2	357	78.0	1913	418.1
TOTAL	193	6.4	2606	86.4	11,001	364.9

* Rate per 100,000 population

† Annual estimated cases and rates by age, gender, race/ethnicity and region are obtained by applying the proportion in the known cases to the total case count.