













California – Chlamydia, Gonorrhea, and P&S Syphilis Rates by County, 2007



Note: Rates are per 100,000 population.
Source: California Department of Public Health, STD Control Branch

California – Chlamydia, Gonorrhea, and P&S Syphilis Rates by Age Group (2007), Race/Ethnicity (2007), and Year



■ Female ■ Male

— Female - - - Male



* Race data may be missing for a substantial number of cases. See the 'Data Limitations' page for further information.
 Note: Rates are per 100,000 population.
 Source: California Department of Public Health, STD Control Branch

California State – Chlamydia, Gonorrhea, and Early Syphilis Cases and Rates Tables for 2007

Gender & Age Group	Chlamydia			Gonorrhea			P&S Syphilis			Early Latent Syphilis			Population	
	Cases	Percent ¹	Rate	Cases	Percent ¹	Rate	Cases	Percent ¹	Rate	Cases	Percent ¹	Rate	Number	Percent ¹
STATE TOTAL	142,912	100.0%	378.4	31,192	100.0%	82.6	2,055	100.0%	5.4	1,460	100.0%	3.9	37,810,582	100.0%
Female Total	101,807	71.2%	537.5	14,490	46.5%	76.5	119	5.8%	0.6	179	12.3%	0.9	18,939,596	50.1%
Ages														
0 – 9	33	0.0%	1.3	15	0.1%	0.6	0	0.0%	0.0	0	0.0%	0.0	2,619,517	13.8%
10 – 14	1,130	1.1%	81.1	214	1.5%	15.4	0	0.0%	0.0	0	0.0%	0.0	1,393,013	7.4%
15 – 19	33,303	32.8%	2,314.7	4,861	33.7%	337.9	16	13.4%	1.1	16	9.0%	1.1	1,438,740	7.6%
20 – 24	36,913	36.4%	2,863.0	4,549	31.5%	352.8	17	14.3%	1.3	34	19.1%	2.6	1,289,314	6.8%
25 – 29	16,718	16.5%	1,403.2	2,332	16.2%	195.7	16	13.4%	1.3	38	21.3%	3.2	1,191,444	6.3%
30 – 34	6,918	6.8%	564.2	1,111	7.7%	90.6	20	16.8%	1.6	32	18.0%	2.6	1,226,258	6.5%
35 – 44	4,923	4.9%	176.2	987	6.8%	35.3	25	21.0%	0.9	34	19.1%	1.2	2,794,590	14.8%
45 +	1,496	1.5%	21.4	365	2.5%	5.2	25	21.0%	0.4	24	13.5%	0.3	6,986,720	36.9%
Not Specified	373	0.4%	-	56	0.4%	-	0	0.0%	-	1	0.6%	-		
Male Total	40,579	28.4%	215.0	16,577	53.1%	87.8	1,934	94.1%	10.2	1,278	87.5%	6.8	18,870,986	49.9%
Ages														
0 – 9	25	0.1%	0.9	10	0.1%	0.4	0	0.0%	0.0	0	0.0%	0.0	2,731,544	14.5%
10 – 14	143	0.4%	9.8	50	0.3%	3.4	1	0.1%	0.1	0	0.0%	0.0	1,455,992	7.7%
15 – 19	8,077	20.0%	532.6	2,292	13.9%	151.1	44	2.3%	2.9	41	3.2%	2.7	1,516,407	8.0%
20 – 24	13,419	33.2%	960.5	4,259	25.8%	304.8	228	11.8%	16.3	146	11.4%	10.5	1,397,128	7.4%
25 – 29	8,441	20.9%	651.4	3,218	19.5%	248.3	316	16.4%	24.4	181	14.2%	14.0	1,295,894	6.9%
30 – 34	3,979	9.9%	310.5	2,058	12.5%	160.6	234	12.1%	18.3	184	14.4%	14.4	1,281,685	6.8%
35 – 44	4,308	10.7%	148.6	3,013	18.2%	103.9	674	34.9%	23.2	424	33.2%	14.6	2,899,150	15.4%
45 +	1,979	4.9%	31.4	1,615	9.8%	25.7	435	22.5%	6.9	301	23.6%	4.8	6,293,186	33.3%
Not Specified	208	0.5%	-	62	0.4%	-	2	0.1%	-	1	0.1%	-		

Gender & Race/Ethnicity	Chlamydia			Gonorrhea			P&S Syphilis			Early Latent Syphilis			Population	
	Cases	Percent ¹	Rate	Cases	Percent ¹	Rate	Cases	Percent ¹	Rate	Cases	Percent ¹	Rate	Number	Percent ¹
STATE TOTAL	142,912	100.0%	378.4	31,192	100.0%	82.6	2,055	100.0%	5.4	1,460	100.0%	3.9	37,810,582	100.0%
Female Total	101,807	71.2%	537.5	14,490	46.5%	76.5	119	5.8%	0.6	179	12.3%	0.9	18,939,596	50.1%
American Indian/Alaska Native	363	0.5%	317.5	64	0.7%	56.0	0	0.0%	0.0	0	0.0%	0.0	114,339	0.6%
Asian/Pacific Islander	4,068	6.1%	171.4	355	3.7%	15.0	1	0.9%	0.0	7	4.0%	0.3	2,372,911	12.5%
Black/African-American	14,381	21.5%	1,243.5	4,241	44.6%	366.7	34	29.1%	2.9	48	27.3%	4.2	1,156,509	6.1%
Latina/Hispanic	34,821	52.0%	526.3	2,927	30.7%	44.2	48	41.0%	0.7	92	52.3%	1.4	6,616,366	34.9%
White	13,298	19.9%	160.6	1,932	20.3%	23.3	34	29.1%	0.4	29	16.5%	0.4	8,278,685	43.7%
Other/Multi/Not Specified	34,876	34.3%	-	4,971	34.3%	-	2	1.7%	-	3	1.7%	-	400,786	2.1%
Male Total	40,579	28.4%	215.0	16,577	53.1%	87.8	1,934	94.1%	10.2	1,278	87.5%	6.8	18,870,986	49.9%
American Indian/Alaska Native	105	0.4%	94.9	48	0.4%	43.4	8	0.4%	7.2	6	0.5%	5.4	110,588	0.6%
Asian/Pacific Islander	1,382	5.2%	63.0	394	3.5%	18.0	76	4.1%	3.5	56	4.5%	2.6	2,193,619	11.6%
Black/African-American	7,301	27.3%	659.4	4,136	37.1%	373.6	255	13.6%	23.0	190	15.3%	17.2	1,107,181	5.9%
Latina/Hispanic	11,919	44.6%	172.1	3,325	29.8%	48.0	643	34.3%	9.3	546	43.9%	7.9	6,923,624	36.7%
White	6,045	22.6%	74.2	3,250	29.1%	39.9	890	47.5%	10.9	445	35.8%	5.5	8,144,845	43.2%
Other/Multi/Not Specified	13,827	34.1%	-	5,424	32.7%	-	62	3.2%	-	35	2.7%	-	391,129	2.1%

¹ Gender specific age groups and race/ethnicity percent calculations exclude "Not Specified" from the denominator.

² Overall state rates were calculated using the July 1 county population estimates from the Department of Finance.
(For California, the July 1, 2007 estimate is 37,771,431).

Note: Rates are per 100,000 population.




Source: California Department of Public Health, STD Control Branch



California – Chlamydia, Gonorrhea, and P&S Syphilis

Ranking of County Rates for 2007

(with 95% Confidence Intervals*)



* Confidence intervals were calculated using Poisson exact method; not shown for counties with zero cases.

Note: Rates are per 100,000 population.

Source: California Department of Public Health, STD Control Branch

