

TABLE 4-10. LEADING CAUSES OF INFANT DEATH BY RACE/ETHNIC GROUP\* OF CHILD, CALIFORNIA, 2001 (By Place of Residence)

RANK	CAUSE OF DEATH	ICD CODES REV. 10	TOTAL INFANT DEATHS	HISPANIC	NON-HISPANIC RACE GROUPS						
					2 OR MORE RACE GROUPS	AMERICAN INDIAN	ASIAN	BLACK	PACIFIC ISLANDER	WHITE/ OTHER RACES	UNKNOWN
	All Causes	A00-Y89	2,815	1,313	54	14	211	390	15	813	5
1	Congenital Malformations, Deformations, and Chromosomal Abnormalities	Q00-Q99	712	385	14	2	55	58	4	193	1
2	Disorders of Short Gestation and Low Birthweight	PO7	371	150	6	2	25	84	1	101	2
3	Sudden Infant Death Syndrome	R95	197	66	10	2	11	31	3	74	-
4	Respiratory Distress of Newborn	P22	98	49	1	-	7	13	1	27	-
4	Complications of Placenta, Cord, and Membrane	P02	98	46	1	2	4	8	1	36	-
6	Effect of Maternal Complications of Pregnancy	P01	87	33	1	-	10	12	-	31	-
7	Bacterial Sepsis of Newborn	P36	71	33	1	-	-	16	-	21	-
8	Neonatal Hemorrhage	P50-P52, P54	62	28	-	-	5	10	-	19	-
9	Intrauterine Hypoxia & Birth Asphyxia	P20-P21	60	31	1	1	3	1	1	22	-
10	Diseases of Circulatory System	I00-I99	58	28	2	-	5	5	-	18	-
11	Accidents	V01-X59, Y85-Y86	53	20	-	-	3	6	1	23	-
	All Other Causes	RESIDUAL	948	444	17	5	83	146	3	248	2

\* Beginning in 2000, up to three races were reported on death certificates. The race/ethnic groups on this table provide a mutually exclusive list with Hispanic, 2 or more race groups, and single races. American Indian includes Eskimo and Aleut, Pacific Islander includes Hawaiian. These groups are only comparable to the year 2000 data tables. See Chapter 6.

- Represents zero events.

Source: State of California, Department of Health Services, Birth and Death Records.